References

1. L.H. Sperling, Polym. Prepr., Am. Chem.Soc., Div. Polym.Chem., 14(2), 958 (1973)
2. M. Matzner, A. Noshay, D.L. Schober and J. E. McGrath, Ind. Cim. Belge., 38, 1104 (1973).
3. J.E. McGrath, L. M. Robeson and M. Matzner, Polym. Prepr., Am. Chem Soc., Div. Polym. Chem., 14(2), 1032 (1973).
4. E.P. Cizek, US Patent 3, 383,435, General Electric Co., 1968.
5. H.E. Bair, Polym.Eng.Sci., 10(4), 247 (1970).
6. N.W. Johnston, Polym.Prepr., Am Chem.Soc., Div. Polym. Chem., 14 (1),46 (1973).
7. Sanghi R, Bhatttacharya B, Singh V. Cassia angustifolia seed gum as an effective natural coagulant for decolourisation of dye solutions. Green Chem. 2002; 4: 252–254
8. Namasivayam C, Arasi DJSE. Removal of congo red from wastewater by adsorption onto waste red mud. Chemosphere 1997; 34: 401–417
9. Moozyckine AU, Davies DM. Green S as a prototype for an environmentally-degradable dye: the concept of a green dye in future Green Chemistry. Green Chem. 2002; 4: 452–458

10. Akhtar S, Khan AA, Husain Q. Potential of immobilized bitter gourd (Momordica charantia) preoxidases in the decolourization and removal of textile dyes from polluted wastewater and dyeing effluent. Chemosphere 2005; 60: 291–301.

11. Crini G. Recent developments in polysaccharide-based materials used as adsorbents in wastewater treatment. Prog. Polym. Sci. 2005; 30: 38–70.

12. Yenkal BF, Yavuz E. Preparation of poly (vinylpyrrolidone) grafted sulfonamide based polystyrene resin and its use for the removal of dye from water. Polym. Adv. Tech. 2006; 17: 928–931.

13. Hashem A, Abdel-Halim ES, Socker HH. Bi-functional starch composites prepared by gamma-irradiation for removal of anionic and cationic dyes from aqueous solutions. Polym. Plast. Tech. Eng. 2007; 46: 71–77.

14. Kasgoz H. Aminofunctionalized acrylamide-maleic acid hydrogels: adsorption of indigo carmine. Coll. Surf. A: Phys. Eng. Asp. 2005; 266: 44–50.

15. Kasgoz H. New sorbent hydrogels for removal of acidic dyes and metal ions from aqueous solutions. Polym. Bull. 2006; 56: 517–528.

16. Dadhaniya PV, Patel MP, Patel RG. Removal of anionic dyes from aqueous solution using poly (N-vinyl pyrrolidone/ 2-(methacryloyloxyethyl)trimetyl ammonium chloride) superswelling hydrogels. Polym. Bull. 2007; 58: 359–369

17. Ohmine I, Tanaka T. Salt effects on the phase transition of ionic gels. J. Chem. Phys. 1982; 77: 5725–5729, Katayama S, Hirokawa Y, Tanaka T.

18. Reentrant phase transition in acrylamide-derivative copolymer gels. Macromolecules 1984; 17: 2641–2643.

19. Zhang LM, Zhou YJ, Wang Y. Novel hydrogel composite for the removal of water-soluble cationic dye. J. Chem. Tech. Biotech. 2006; 81: 799–804k: preparation, swelling and dye adsorption properties. Polym.